

European Journal of Inorganic Chemistry 2014 vol.2014 N34, pages 5958-5968

σ 2P,O-hybrid ligands: Synthesis of the first 4-hydroxy-1,3-benzazaphospholes by ortho-lithiation of m-amidophenyl diethyl phosphates

Aluri B., Shah K., Gupta N., Fomina O., Yakhvarov D., Ghalib M., Jones P., Schulzke C., Heinicke J.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

Copyright © 2014 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim. The m-phosphorylanilides 2 are available from anilides 1 by the Atherton-Todd reaction; the selective ortho-lithiation of the o'-methyl-protected phosphorylpivalanilide 2b with tBuLi proceeded in high yield in the presence of ClSiMe₃. The ortho-lithiation is followed by rapid 1,3-migration of the PO₃Et₂ group to yield the phosphonoanilide cis/trans-3b. This compound mainly reacts with excess LiAlH₄ by reductive cyclization to form the 4-hydroxy-1H-1,3-benzazaphosphole 6. The lithiation of the o'-unprotected phosphorylpivalanilide 2a with LDA was unselective and led to 3a and 4a in low yields, whereas additional ortho-lithiation of the benzoyl group occurred for the lithiation of the o'-protected phosphonobenzanilide 2c with tBuLi/LDA to give 7 in rather low yield. The reduction of crude 7 led to (benzylamino)phenol 8 and the 4-hydroxy-1H-1,3-benzazaphosphole 9 as a minor product. The properties, NMR spectroscopy data, and crystal structures of 5b, 6, and 8 are reported.

<http://dx.doi.org/10.1002/ejic.201402527>

Keywords

Amides, Lithiation, Phosphorus heterocycles, Phosphorylation, Rearrangement